

A DEFORMATION OF THE ORLIK-SOLOMON ALGEBRA

ISTVÁN HECKENBERGER AND VOLKMAR WELKER

ABSTRACT. A deformation of the Orlik-Solomon algebra of a matroid $\mathfrak{M}$ is defined as a quotient of the free associative algebra over a commutative ring R with 1. It is shown that the given generators form a Gröbner basis and that after suitable homogenization the deformation and the Orlik-Solomon have the same Hilbert series as R -algebras. For supersolvable matroids, equivalently fiber type arrangements, there is a quadratic Gröbner basis and hence the algebra is Koszul.

1. INTRODUCTION AND STATEMENT OF RESULTS

In this paper we introduce and study a deformation of the Orlik-Solomon algebra of a matroid $\mathfrak{M}$. We refer the reader to [Orl07] and [Yuz01] for general facts about the classical Orlik-Solomon algebra. Our deformation, which is different from the one in [SY97], is presented as a quotient of the free associative algebra over some commutative ring R with 1 by an ideal $I_q(\mathfrak{M})$ whose generators are deformations of the classical generators of the defining ideal of the Orlik-Solomon algebra by a parameter $q \in R$. Choosing $q = 0$ yields the Orlik-Solomon algebra over R . Our main result, Theorem 1.1, states that the given generators of $I_q(\mathfrak{M})$ are a Gröbner basis of the ideal. As a consequence it is shown in Corollary 1.3 that the deformation with q regarded as a degree 2 element is a standard graded R -algebra which has the same Hilbert series as the Orlik-Solomon algebra. For supersolvable matroids, equivalently fiber type arrangements, the existence of a quadratic Gröbner basis is shown which implies that the algebra is Koszul. As further consequences we obtain in Corollary 1.5 a known Gröbner basis for the Orlik-Solomon algebra as a quotient of the free and the exterior algebra. The remaining part of the introduction is devoted to the basic definitions and statement of results. In Section 2 basic facts about non-commutative Gröbner basis theory are given. Section 3 provides technical lemmas needed for the proof of the main result. Finally in Section 4 the missing proofs are given and an independence statement is presented.

Let S be a finite set and fix a total order $<$ on S . Let R be a commutative ring with unit 1 and let $q \in R$. For an arbitrary set system $\mathfrak{M} \subseteq 2^S$ we define a two-sided ideal $I_q(\mathfrak{M})$ in the ring

$$A := R\langle t_s \mid s \in S \rangle$$

of non-commutative polynomials in the variables t_s , $s \in S$, with coefficients in R .

2000 *Mathematics Subject Classification.* Primary: 05B35 Secondary: 16S37, 16S80.

Let $\mathcal{I}_q(\mathfrak{M})$ denote the subset of A consisting of the elements

$$(1.1) \quad t_s^2 - q, \quad s \in S,$$

$$(1.2) \quad t_r t_s + t_s t_r - 2q, \quad r, s \in S, s < r,$$

$$(1.3) \quad t_J^- := \sum_{I \subseteq J, 2 \nmid \#I} (-1)^{\ell_J(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}, \quad \text{for all } J \in \mathfrak{M}.$$

Here, for a subset $I = \{j_{\alpha_1} < \cdots < j_{\alpha_{\#I}}\} \subseteq J = \{j_1 < \cdots < j_{\#J}\}$ we set

$$\ell_J(I) = \sum_{\nu=1}^{\#I} (\alpha_\nu - \nu).$$

The two-sided ideal of A generated by $\mathcal{I}_q(\mathfrak{M})$ will be denoted by $I_q(\mathfrak{M})$. We write $\text{OS}_q(\mathfrak{M})$ for the quotient $A/I_q(\mathfrak{M})$. Our main motivation comes from the situation when $q = 0$ and $\mathfrak{M}$ is indeed the set of circuits of a loopless matroid without parallel elements. We refer the reader to the books [Wel76] and [Oxl06] as a general reference for matroid theory and recall that for loopless matroids without parallel element set of circuits $\mathfrak{M}$ is characterized by the following three axioms:

(C1) $J \in \mathfrak{M} \Rightarrow \#J > 2$.

(C2) $J, K \in \mathfrak{M}, J \subseteq K \Rightarrow J = K$.

(C3) For any $J, K \in \mathfrak{M}$ such that $J \neq K$ and for any $x \in J \cap K$ there exists $L \in \mathfrak{M}$ such that $L \subseteq (J \cup K) \setminus \{x\}$.

We will refer to (C3) also by the name *circuit axiom*.

Now if $\mathfrak{M}$ is the set of circuits of a loopless matroid without parallel elements then for $q = 0$ the algebra $\text{OS}_q(\mathfrak{M})$ is the *Orlik-Solomon algebra* of $\mathfrak{M}$. In case $\mathfrak{M}$ is realizable as the set of circuits of a finite set of hyperplanes in $\mathbb{C}^d$ then the Orlik-Solomon algebra of $\mathfrak{M}$ is known to be the cohomology algebra of the set-theoretic complement of the union of the hyperplanes [OS80, (5.2)]. In general, for $q \neq 0$ the algebra $\text{OS}_q(\mathfrak{M})$ is not isomorphic to the Orlik-Solomon algebra of $\mathfrak{M}$, take for example S with $\#S = 1$. On the other hand, if q is a square in R , then $\text{OS}_q(\mathfrak{M})$ is easily seen to be isomorphic to $\text{OS}_1(\mathfrak{M})$.

Before we can proceed to the statement of our main results we need some more definitions. Throughout this paper we will use the *degree lexicographic* order $<_{\text{dlex}}$ (see Section 2) on the monomials in A induced by the total order $<$ on S as a term order for the monomials in A . We enumerate the elements of any subset $J \subseteq S$ by $j_1, \dots, j_{\#J}$ such that $j_1 < \cdots < j_{\#J}$. For any $J \subseteq S$ we write t_J for the monomial $t_{j_1} \cdots t_{j_{\#J}} \in A$. The degree lexicographic order on the monomials t_J induces the degree lexicographic order on subsets of S ; that is for subsets $J, K \subseteq S$ we set $J <_{\text{dlex}} K$ if and only if $\#J < \#K$ or $\#J = \#K$ and the minimum of the symmetric difference of J and K is contained in J . In addition, for two subsets $K, J \subseteq S$ we say that K is a *convex subset* of J if $K \subseteq J$ and if $j \in K$ for all $j \in J$ with $k < j < k'$ for some $k, k' \in K$. We write $K \sqsubseteq J$ in this situation. Recall that a subset $J \subseteq S$ is called *dependent* in $\mathfrak{M}$ if it contains a circuit from $\mathfrak{M}$. Let $\overline{\mathfrak{M}}$ be the set of dependent subsets of S defined recursively as follows:

(GC) A dependent set $J \subseteq S$ belongs to $\overline{\mathfrak{M}}$ if and only if $K \in \overline{\mathfrak{M}}$ with $K <_{\text{dlex}} J$ implies that $K \setminus \{k_1\} \not\subseteq J \setminus \{j_1\}$.

Then we call $\overline{\mathfrak{M}}$ the set of *Gröbner circuits* of $\mathfrak{M}$.

Theorem 1.1. *Let $\overline{\mathfrak{M}}$ be a set of Gröbner circuits of a loopless matroid $\mathfrak{M}$ without parallel elements. Then $I_q(\mathfrak{M}) = I_q(\overline{\mathfrak{M}})$ and the set $\mathcal{I}_q(\overline{\mathfrak{M}})$ is a Gröbner basis of the ideal $I_q(\mathfrak{M})$ with respect to the degree lexicographic order for all choices of q .*

The Gröbner basis is easily seen to depend on the total order chosen on S but in Proposition 4.1 we show that $I_q(\mathfrak{M})$ and hence $\text{OS}_q(\mathfrak{M})$ is independent of the order on S . Simple inspection shows that the leading monomial of (1.1) is t_s^2 , of (1.2) is $t_r t_s$ for $s < r$ and of (1.3) is $t_{J \setminus \{j_1\}}$. Thus all leading coefficients of $\mathcal{I}_q(\overline{\mathfrak{M}})$ are 1. The standard monomials with respect to the Gröbner basis $\mathcal{I}_q(\overline{\mathfrak{M}})$ are the monomials $m = t_J$ for $J \subseteq S$ for which there is no factorization $m = m_1 t_{K'} m_2$ for monomials m_1, m_2 where K' is a *broken Gröbner circuit*; that is there is a Gröbner circuit K for which $K' = K \setminus \{k_1\}$. By the definition of a Gröbner circuit it then follows that the standard monomials with respect to the Gröbner basis $\mathcal{I}_q(\overline{\mathfrak{M}})$ are the monomials $m = t_J$ such that J does not contain a *broken circuit* of $\mathfrak{M}$; that is a circuit with its least element removed. These facts immediately imply:

Corollary 1.2. *Let $\mathfrak{M}$ be a loopless matroid without parallel elements. Then the algebra $\text{OS}_q(\mathfrak{M})$ is a free R -module whose rank is independent of q .*

The standard monomials with respect to the Gröbner basis $\mathcal{I}_q(\overline{\mathfrak{M}})$ are the monomials $m = t_J$ for $J \subseteq S$ for which J does not contain a broken circuit.

The algebra $\text{OS}_q(\mathfrak{M})$ is $\mathbb{Z}/2\mathbb{Z}$ -graded for the grading induced by $\deg t_s = 1$ for all $s \in S$. If R is $\mathbb{Z}$ -graded and $t_0 \in R$ is homogeneous of degree one then the $\mathbb{Z}$ -grading of R extends to a $\mathbb{Z}$ -grading of $\text{OS}_q(\mathfrak{M})$ for $q = t_0^2$. We consider the case $R = Q[t_0]$ for some commutative ring Q with 1 and extend the total order on the variables by setting t_0 to be the least variable. Then we consider $I_q(\mathfrak{M})$ as an ideal in $Q\langle t_s | s \in S \rangle[t_0]$. We deduce from Theorem 1.1, standard facts about homogenizing Gröbner bases (see [Li02, Thm. 3.7]) and Corollary 1.2 the following corollary.

Corollary 1.3. *Let $\mathfrak{M}$ be a loopless matroid without parallel elements, $R = Q[t_0]$ for a commutative ring Q with 1 and t_0 a degree one variable. If we set $q = t_0^2$ then*

- (1) *the set $\mathcal{I}_q(\overline{\mathfrak{M}})$ is a Gröbner basis of $I_q(\mathfrak{M})$ for the degree lexicographic order with t_0 being the least variable. The algebra $\text{OS}_q(\mathfrak{M})$ is a free Q -module and a standard graded Q -algebra.*
- (2) *The Hilbert series of $\text{OS}_q(\mathfrak{M})$ as a Q -algebra is*

$$\frac{1 + c_1 z + \cdots + c_{\text{rk}(\mathfrak{M})} z^{\text{rk}(\mathfrak{M})}}{1 - z} = \text{Hilb}_{\mathfrak{M}}(z) \cdot \frac{1}{1 - z},$$

where $\text{rk}(\mathfrak{M})$ is the rank of $\mathfrak{M}$, c_i is the number of subsets of S of cardinality i not containing a broken circuit and $\text{Hilb}_{\mathfrak{M}}(z)$ the Hilbert series of the Orlik-Solomon algebra of $\mathfrak{M}$ as an algebra over a field k .

Note that the first part together with the second part of Corollary 1.2 implies that the standard monomials of the Gröbner basis are $q^i t_J$ for $i \geq 0$ and $J \subset S$ such that J does not contain a broken circuit. Part (2) of Corollary 1.3 hence follows by a simple counting argument and standard facts about Orlik-Solomon algebras.

We note that experiments suggest that the generators (1.1), (1.2) and (1.3) are the unique deformations of the corresponding polynomials for $q = 0$ by variables of degree ≥ 1 satisfying Corollary 1.3(2).

Using results from matroid theory [BZ91] we obtain the following results extending results from [SY97] (Koszul property) and [Pee03] (quadratic Gröbner basis and Koszul property) for Orlik-Solomon algebras to our deformation. We refer the reader to [Frö99] for basic facts about Koszul algebras.

Corollary 1.4. *Let $\mathfrak{M}$ be a supersolvable loopless matroid without parallel elements, $R = Q[t_0]$ for a commutative ring Q with 1 and t_0 a degree one variable. If we set $q = t_0^2$ then $\mathcal{I}_q(\overline{\mathfrak{M}})$ is a quadratic Gröbner basis and in particular $\text{OS}_q(\mathfrak{M})$ is a standard graded Koszul algebra.*

We postpone the derivation of this corollary till Section 4. For $q = 0$, Theorem 1.1 states

Corollary 1.5. *Let $\overline{\mathfrak{M}}$ be the set of Gröbner circuits of a loopless matroid $\mathfrak{M}$ without parallel elements. Then the polynomials t_s^2 with $s \in S$, $t_r t_s + t_s t_r$ with $r, s \in S$, $s < r$ and $\sum_{\nu=1}^{\#J} (-1)^{\nu-1} t_{J \setminus \{j_\nu\}}$, where $J \in \overline{\mathfrak{M}}$, form a Gröbner basis of the defining ideal $I_0(\mathfrak{M})$ of the Orlik-Solomon algebra of $\mathfrak{M}$ with respect to the degree lexicographic order.*

Since for $q = 0$ the quotient of A by t_s^2 with $s \in S$, $t_r t_s + t_s t_r$ with $r, s \in S$, $s < r$ is the exterior algebra E we also get the following corollary from [Mor94, Prop. 9.3]. For its formulation we identify $t_J \in A$ for $J \subseteq S$ with its image in E .

Corollary 1.6. *Let $\overline{\mathfrak{M}}$ be the set of Gröbner circuits of a loopless matroid $\mathfrak{M}$ without parallel elements. Then the polynomials $\sum_{\nu=1}^{\#J} (-1)^{\nu-1} t_{J \setminus \{j_\nu\}}$, where $J \in \mathfrak{M} \cap \overline{\mathfrak{M}}$, form a Gröbner basis of the defining ideal of the Orlik-Solomon algebra in E .*

Gröbner bases of the defining ideal of the Orlik-Solomon algebra inside the exterior algebra have been described previously (see for example [Yuz01, Thm. 2.8], [CF05], [Pee03]).

2. NON-COMMUTATIVE GRÖBNER BASICS

Recall that the *degree lexicographic order* or *deglex order* on the monomials in A is the total order $<_{\text{dlex}}$ such that for two monomials $t_{i_1} \cdots t_{i_k}$ and $t_{j_1} \cdots t_{j_l}$ in A we have $t_{i_1} \cdots t_{i_k} <_{\text{dlex}} t_{j_1} \cdots t_{j_l}$ if and only if either $k < l$ or $k = l$ and for some $0 \leq h < k$ we have $i_1 = j_1, \dots, i_h = j_h$ and $i_{h+1} < j_{h+1}$. Any $\xi \in A$ can uniquely be written as a polynomial of the form $f = c_1 m_1 + \cdots + c_j m_j$, for non-commutative monomials $m_j <_{\text{dlex}} \cdots m_2 <_{\text{dlex}} m_1$ in the variables t_s , $s \in S$, and ring elements $c_1, \dots, c_j \in R \setminus \{0\}$. In this polynomial, $c_1 m_1$ is called the *leading term*, c_1 the *leading coefficient* and m_1 the *leading monomial* of f . We write $\text{lt}(f)$ for the leading term, $\text{lm}(f)$ for the leading monomial and $\text{lc}(f)$ for the leading coefficient of f . The $m_1, \dots, m_k$ are called the *monomials of f* . In other words, the leading

monomial is the largest monomial among all monomials of f with respect to the deglex order. Further, for any monomial $m \in A$ there are only finitely many monomials $m' \in A$ such that $m' <_{\text{dlex}} m$.

Let $\mathcal{I}$ be a set of elements of A with leading coefficient 1. A *reduction* of a polynomial $f \in A$ modulo $\mathcal{I}$ is an expression obtained from f by replacing the leading monomial m of an element $g \in \mathcal{I}$, appearing as a subword of one of the monomials of f , by $m - g$. By construction, a reduction does not have monomials larger than the leading monomial of f . For any $f, g \in A$ we say that f *reduces to* g (modulo $\mathcal{I}$) and write

$$(2.1) \quad f \searrow_{\mathcal{I}} g$$

if there is a sequence of expressions $f = f_0, f_1, \dots, f_k = g$, where $k \in \mathbb{N}_0$, such that f_{i+1} is a reduction of f_i for all $i \in \{0, 1, \dots, k-1\}$.

A subset G of a two-sided ideal I in A is called a *Gröbner basis* of I if the two-sided ideal generated by $\{\text{lt}(g) \mid g \in G\}$ coincides with the two-sided ideal generated by $\{\text{lt}(f) \mid f \in I\}$.

For two polynomials f, g in A with $\text{lc}(f) = \text{lc}(g) = 1$ an *S-polynomial* of (f, g) is any non-zero expression $m_1 f m_2 - n_1 g n_2 \in A$ for monomials m_1, m_2, n_1, n_2 such that

$$(2.2) \quad m_1 \text{lm}(f) m_2 = n_1 \text{lm}(g) n_2.$$

Let $J_{f,g}$ be the submodule of the A -bimodule $(A \oplus A) \otimes_R (A \oplus A)$ generated by the tensors $(m_1, n_1) \otimes (m_2, n_2)$ for monomials m_1, m_2, n_1, n_2 for which (2.2) holds. Being generated by tensors of pairs of monomials there is a unique inclusionwise minimal set of generators of $J_{f,g}$ consisting of tensors of pairs of monomials. It is easily seen that any of the generators will be of the form $(m, 1) \otimes (1, n)$, $(1, n) \otimes (m, 1)$, $(1, n_1) \otimes (1, n_2)$ or $(m_1, 1) \otimes (m_2, 1)$. The criterion from the following theorem will be employed in order to derive Theorem 1.1.

Theorem 2.1. *Let R be a field and I a two-sided ideal of A . A set $\mathcal{I} := \{f_1, \dots, f_r\} \subseteq I$ is a Gröbner basis for I if and only if for all $1 \leq i \leq j \leq r$ and for any minimal generator $(m_1, n_1) \otimes (m_2, n_2)$ of J_{f_i, f_j} the corresponding S-polynomial of (f_i, f_j) reduces to 0 modulo $\mathcal{I}$.*

It is possible to simplify the Gröbner basis criterion in Theorem 2.1 by using the following fact [Mor94, Cor. 5.8].

Lemma 2.2. *Let $f, g \in \mathcal{I}$. Then the S-polynomials of (f, g) corresponding to the generators $(\text{lm}(g) m, 1) \otimes (1, m \text{lm}(f))$ and $(1, \text{lm}(f) m) \otimes (m \text{lm}(g), 1)$ of $J_{f,g}$, where m is an arbitrary monomial, reduce to 0 modulo $\mathcal{I}$.*

We will apply Theorem 2.1 and Lemma 2.2 in a situation where R is not necessarily a field. But since all our polynomials have leading coefficient 1 and since all reductions only use coefficients ± 1 the assertions remain valid.

3. TECHNICAL LEMMAS

3.1. General set systems. In this section we collect some useful formulas which are valid for arbitrary set systems $\mathfrak{M}$ over S .

Generalizing the notation in the introduction, for all $J \subseteq S$ let

$$t_J^- = \sum_{I \subseteq J, 2 \nmid \#I} (-1)^{\ell_J(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}, \quad t_J^+ = \sum_{I \subseteq J, 2 \mid \#I} (-1)^{\ell_J(I)} (-q)^{\#I/2} t_{J \setminus I}.$$

We start with deriving formulas which are valid in A .

Lemma 3.1. *Let $J, J', J'' \subseteq S$ such that $J = J' \cup J''$ and $j' < j''$ for all $j' \in J', j'' \in J''$. Then in A equations*

$$(3.1) \quad t_J^+ = t_{J'}^+ t_{J''}^+ + (-1)^{\#J'} q t_{J'}^- t_{J''}^-,$$

$$(3.2) \quad t_J^- = t_{J'}^- t_{J''}^+ + (-1)^{\#J'} t_{J'}^+ t_{J''}^-,$$

hold.

Proof. We proceed by induction on $\#J'$. Since $t_\emptyset^+ = 1$ and $t_\emptyset^- = 0$, the claim holds for $J' = \emptyset$. Assume now that $\#J \geq 1$, $J' = \{j_1\}$ and $J'' = J \setminus \{j_1\}$. Then

$$\begin{aligned} t_J^+ &= \sum_{I \subseteq J, 2 \nmid \#I} (-1)^{\ell_J(I)} (-q)^{\#I/2} t_{J \setminus I} \\ &= \sum_{I \subseteq J, 2 \mid \#I, j_1 \in I} (-1)^{\ell_J(I)} (-q)^{\#I/2} t_{J \setminus I} + \sum_{I \subseteq J, 2 \mid \#I, j_1 \notin I} (-1)^{\ell_J(I)} (-q)^{\#I/2} t_{J \setminus I} \\ &= \sum_{L \subseteq J'', 2 \nmid \#L} (-1)^{\ell_{J''}(L)} (-q)^{(\#L+1)/2} t_{J'' \setminus L} + \sum_{L \subseteq J'', 2 \mid \#L} (-1)^{\ell_{J''}(L) + \#L} (-q)^{\#L/2} t_{j_1} t_{J'' \setminus L} \\ &= -q t_{J''}^- + t_{j_1} t_{J''}^+ = t_{J'}^+ t_{J''}^+ - q t_{J'}^- t_{J''}^-. \end{aligned}$$

Similarly,

$$t_J^- = t_{J''}^+ - t_{j_1} t_{J''}^- = t_{J'}^- t_{J''}^+ - t_{J'}^+ t_{J''}^-.$$

The rest follows from the induction hypothesis and the associativity law of A . $\square$

Lemma 3.2. *Let $K, L \subseteq S$ such that $k < l$ for all $k \in K, l \in L$. Then in A we have*

$$(3.3) \quad t_K^- t_L^- = \sum_{i=1}^{\#K} (-1)^{\#K-i} t_{L \cup K \setminus \{k_i\}}^- = \sum_{i=1}^{\#L} (-1)^{i-1} t_{K \cup L \setminus \{l_i\}}^-.$$

Proof. We prove the first equality by induction on $\#K$. If $K = \emptyset$ then both sides of the equality are zero. Assume now that $K \neq \emptyset$ and let $n = \#K$, $K' = K \setminus \{k_n\}$ and $L' = \{k_n\} \cup L$. Then Lemma 3.1 applied three times implies that

$$\begin{aligned} t_K^- t_L^- &= (t_{K'}^- t_{k_n} + (-1)^{n-1} t_{K'}^+ t_L^-) t_L^- \\ &= t_{K'}^- (-t_{L'}^- + t_L^+) + (-1)^{n-1} t_{K'}^+ t_{K'}^- t_L^- \\ &= -t_{K'}^- t_{L'}^- + t_{L \cup K \setminus \{k_n\}}^- \end{aligned}$$

from which the first equation follows from the induction hypothesis. In particular, for $L = \emptyset$ we obtain that $0 = \sum_{i=1}^{\#J} (-1)^{\#J-i} t_{J \setminus \{j_i\}}^-$ for all $J \subseteq S$. This implies that the second and the third expression in (3.3) coincide. $\square$

Lemma 3.3. *Let $J, L \subseteq S$ such that $n := \#J \geq 2$, $J \setminus \{j_1, j_n\} = L \setminus \{l_1\}$ and $j_1 < l_1$. Then in A we have*

$$(3.4) \quad t_J^- - t_L^- t_{j_n} - \sum_{i=2}^{n-1} (-1)^i t_{(J \cup L) \setminus \{j_i\}}^- + (-1)^n t_{j_1} t_L^- = 0.$$

Proof. Let $J' = J \setminus \{j_1\}$ and $L' = L \setminus \{l_1\}$. Then $t_J^- = t_{J'}^+ - t_{j_1} t_{J'}^-$ by Lemma 3.1 and

$$\sum_{i=2}^{n-1} (-1)^i t_{(J \cup L) \setminus \{j_i\}}^- = (t_{l_1} - t_{j_1}) t_{J'}^- - (-1)^n t_{(J \cup L) \setminus \{j_n\}}^-$$

by Lemma 3.2. Thus Equation (3.4) is equivalent to

$$t_{J'}^+ - t_L^- t_{j_n} - t_{l_1} t_{J'}^- + (-1)^n t_{(J \cup L) \setminus \{j_n\}}^- + (-1)^n t_{j_1} t_L^- = 0.$$

By Lemma 3.1, the left hand side of the latter equation can be written as

$$\begin{aligned} & (t_{L'}^+ t_{j_n} + (-1)^{n-2} q t_{L'}^-) - (t_{L'}^+ - t_{l_1} t_{L'}^-) t_{j_n} - t_{l_1} (t_{L'}^- t_{j_n} + (-1)^{n-2} t_{L'}^+) \\ & + (-1)^n (t_L^+ - t_{j_1} t_L^-) + (-1)^n t_{j_1} t_L^- \\ & = (-1)^{n-2} q t_{L'}^- + (-1)^{n-1} t_{l_1} t_{L'}^+ + (-1)^n t_L^+ = 0 \end{aligned}$$

which proves the claim. $\square$

Lemma 3.4. *Let $J, L \subseteq S$ such that $n := \#J \geq 2$, $L \setminus \{l_1\} = J \setminus \{j_1, j_2\}$ and $j_1 < l_1 < j_2$. Then in A we have*

$$(3.5) \quad t_J^- - (t_{j_2} t_{l_1} + t_{l_1} t_{j_2} - 2q) t_{L \setminus \{l_1\}}^- - t_{j_1 j_2}^- t_L^- + \sum_{i=3}^n (-1)^{i+1} t_{(J \cup L) \setminus \{j_i\}}^- = 0.$$

Proof. Let $J' = J \setminus \{j_1\}$ and $L' = L \setminus \{l_1\}$. Then

$$\sum_{i=3}^n (-1)^{i+1} t_{(J \cup L) \setminus \{j_i\}}^- = t_{\{j_1, l_1, j_2\}}^- t_{L'}^-$$

by Lemma 3.2. Applying this and Lemma 3.1 repeatedly, the left hand side of Equation (3.4) becomes

$$\begin{aligned} & t_J^- - (t_{j_2} t_{l_1} + t_{l_1} t_{j_2} - 2q) t_{L'}^- - (t_{j_2} - t_{j_1}) (t_{L'}^+ - t_{l_1} t_{L'}^-) + (t_{l_1} t_{j_2} - t_{j_1} t_{j_2} + t_{j_1} t_{l_1} - q) t_{L'}^- \\ & = t_J^- + (q - t_{j_1} t_{j_2}) t_{L'}^- - (t_{j_2} - t_{j_1}) t_{L'}^+ \\ & = t_{J'}^+ - t_{j_1} t_{J'}^- + (q - t_{j_1} t_{j_2}) t_{L'}^- - (t_{j_2} - t_{j_1}) t_{L'}^+ = 0 \end{aligned}$$

which proves the claim. $\square$

Now we turn to reductions. Observe that if $\mathfrak{N} \subseteq \mathfrak{M}$ are sets of subsets of S then for $t \in A$ we have that if t reduces to zero modulo $\mathcal{I}_q(\mathfrak{N})$ then t reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$. In particular, if t reduces to zero modulo $\mathcal{I}_q(\emptyset)$ then t reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$. Note that in the reduction modulo $\mathcal{I}_q(\emptyset)$ only relations (1.1) and (1.2) are involved.

Lemma 3.5. *Let $J \subseteq S$ and let $s \in S$.*

(1) Assume that $s \in J$. Then

$$t_s t_J^+ \searrow^{\mathcal{I}_q(\emptyset)} q t_J^-, \quad t_s t_J^- \searrow^{\mathcal{I}_q(\emptyset)} t_J^+.$$

(2) Assume that $s \notin J$. Let $J' = \{j \in J \mid j < s\}$. Then

$$t_s t_J^+ \searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J'} t_{J \cup \{s\}}^+ + q t_J^-, \quad t_s t_J^- \searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J'-1} t_{J \cup \{s\}}^- + t_J^+.$$

Remark 3.6. If $s < j_1$ then in the last expression of Lemma 3.5(2) the leading term of $t_s t_J^-$ is $t_s t_{j_2} \cdots t_{j_{\#J}}$. On the other hand, the leading term of both $t_{J \cup \{s\}}^-$ and t_J^+ is t_J which is larger than $t_s t_{j_2} \cdots t_{j_{\#J}}$ with respect to the deglex order. The reduction formula means that these two leading terms cancel and $t_s t_J^-$ reduces modulo $\mathcal{I}_q(\emptyset)$ to the remaining expression. In fact, due to Equation (3.2) for $t_{J \cup \{s\}}^-$, no reduction is needed to obtain the result.

Proof. We proceed by induction on $\#J$. Assume first that $s < j$ for all $j \in J$. (This holds in particular if $J = \emptyset$.) Then $t_{J \cup \{s\}}^+ = t_s t_J^+ - q t_J^-$ and $t_{J \cup \{s\}}^- = t_J^+ - t_s t_J^-$ in A by Lemma 3.1 and hence (2) holds in this case.

Assume now that $s \in J$ and $s \leq j$ for all $j \in J$. Let $K = J \setminus \{s\}$. Then

$$t_J^+ = t_s t_K^+ - q t_K^-, \quad t_J^- = t_K^+ - t_s t_K^-$$

in A by Lemma 3.1. Since $t_s^2 - q \in \mathcal{I}_q(\emptyset)$, it follows that

$$\begin{aligned} t_s t_J^+ &= t_s^2 t_K^+ - q t_s t_K^- \searrow^{\mathcal{I}_q(\emptyset)} q t_K^+ - q t_s t_K^- = q t_J^-, \\ t_s t_J^- &= t_s t_K^+ - t_s^2 t_K^- \searrow^{\mathcal{I}_q(\emptyset)} t_s t_K^+ - q t_K^- = t_J^+ \end{aligned}$$

by Lemma 3.1. Hence (1) holds in this case.

Finally, assume that $J \neq \emptyset$ and that $s > j_1$. Let $K = J \setminus \{j_1\}$. We assume first that $s \in J$ and prove (1). Since $t_s t_{j_1} + t_{j_1} t_s - 2q \in \mathcal{I}_q(\emptyset)$, by Lemma 3.1 and by induction hypothesis we obtain that

$$\begin{aligned} t_s t_J^+ &= t_s(t_{j_1} t_K^+ - q t_K^-) \searrow^{\mathcal{I}_q(\emptyset)} (-t_{j_1} t_s + 2q) t_K^+ - q t_s t_K^- \\ &\searrow^{\mathcal{I}_q(\emptyset)} -t_{j_1}(q t_K^-) + q t_K^+ = q t_J^-. \end{aligned}$$

Similarly,

$$\begin{aligned} t_s t_J^- &= t_s(t_K^+ - t_{j_1} t_K^-) \searrow^{\mathcal{I}_q(\emptyset)} q t_K^- - (-t_{j_1} t_s + 2q) t_K^- \\ &\searrow^{\mathcal{I}_q(\emptyset)} t_{j_1} t_K^+ - q t_K^- = t_J^+. \end{aligned}$$

A similar argument proves (2). □

The following lemma is a right-handed analogue of the previous result.

Lemma 3.7. Let $J \subseteq S$ and let $s \in S$.

(1) Assume that $s \in J$. Then

$$t_J^+ t_s \searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J+1} q t_J^-, \quad t_J^- t_s \searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J+1} t_J^+.$$

(2) Assume that $s \notin J$. Let $J'' = \{j \in J \mid s < j\}$. Then

$$\begin{aligned} t_J^+ t_s &\searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J''} t_{J \cup \{s\}}^+ + (-1)^{\#J+1} q t_J^-, \\ t_J^- t_s &\searrow^{\mathcal{I}_q(\emptyset)} (-1)^{\#J''} t_{J \cup \{s\}}^- + (-1)^{\#J+1} t_J^+. \end{aligned}$$

Proof. See the proof of Lemma 3.5. $\square$

Lemma 3.8. Let $J \subseteq S$ with $J \neq \emptyset$. If t_J^- reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$, then t_J^+ reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$.

Proof. Let $K = J \setminus \{j_1\}$. Lemma 3.1 gives that

$$t_J^+ = t_{j_1} t_K^+ - q t_K^-, \quad t_J^- = t_K^+ - t_{j_1} t_K^-$$

in A and the leading term of t_J^- is the leading term of t_K^+ . Thus, since $t_J^- \searrow^{\mathcal{I}_q(\mathfrak{M})} 0$, it follows that $t_K^+ \searrow^{\mathcal{I}_q(\mathfrak{M})} t_{j_1} t_K^-$. Hence

$$t_J^+ = t_{j_1} t_K^+ - q t_K^- \searrow^{\mathcal{I}_q(\mathfrak{M})} t_{j_1}^2 t_K^- - q t_K^-$$

which reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$ since $t_{j_1}^2 - q \in \mathcal{I}_q(\mathfrak{M})$. $\square$

Lemma 3.9. Let $J \subseteq S$ and let $s \in S$. Assume that t_J^- reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$. If $J \neq \emptyset$, $r < s$ for all $r \in J$ or $\#J \geq 2$, $s < j_2$ then $t_{J \cup \{s\}}^-$ reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$.

Proof. If $r < s$ for all $r \in J$ then $t_{J \cup \{s\}}^- = t_J^- t_s - t_J^+$ by Lemma 3.1. Thus the first half of the claim holds by Lemma 3.8.

If $\#J \geq 2$, $s < j_1$ then $t_{J \cup \{s\}}^- = t_J^+ - t_s t_J^-$ by Lemma 3.1 and again the claim holds. If $s = j_1$ then there is nothing to prove. Finally, if $j_1 < s < j_2$ then let $J' = J \setminus \{j_1\}$. Lemma 3.1 gives that

$$\begin{aligned} t_{J \cup \{s\}}^- &= t_{\{j_1, s\}}^- t_{J'}^+ + t_{\{j_1, s\}}^+ t_{J'}^- = (t_s - t_{j_1}) t_{J'}^+ + (t_{j_1} t_s - q) t_{J'}^-, \\ &= (t_s - t_{j_1}) (t_J^- + t_{j_1} t_{J'}^-) + (t_{j_1} t_s - q) t_{J'}^-, \\ &= (t_s - t_{j_1}) t_J^- + (t_s t_{j_1} + t_{j_1} t_s - 2q) t_{J'}^- - (t_{j_1}^2 - q) t_{J'}^-. \end{aligned}$$

This expression reduces to zero modulo $\mathcal{I}_q(\mathfrak{M})$ which proves the remaining claim. $\square$

3.2. Matroids. From now on let $\mathfrak{M}$ be the set of circuits of a loopless matroid without parallel elements on ground set S and let $\overline{\mathfrak{M}}$ be a set of Gröbner circuits of $\mathfrak{M}$.

Example 3.10. A typical example where the set of circuits $\mathfrak{M}$ of a matroid is not sufficient to define a Gröbner basis of $\text{OS}_q(\mathfrak{M})$ is the following.

Let $S = \{1, 2, 3, 4\}$ with the usual order and let $\mathfrak{M}$ be the set system consisting of $\{1, 2, 4\}$. Then

$$t_{1234}^- = t_2 t_3 t_4 - t_1 t_3 t_4 + t_1 t_2 t_4 - t_1 t_2 t_3 - q t_4 + q t_3 - q t_2 + q t_1$$

is zero in $\text{OS}_q(\mathfrak{M})$ since

$$t_{1234}^- = -t_3 t_{124}^- + t_{124}^+ = -t_3 t_{124}^- + t_1 t_{124}^-$$

by Lemma 3.5 and $t_{124}^- = 0$ in $\text{OS}_q(\mathfrak{M})$. However, the leading term of t_{1234}^- cannot be reduced using the generators of $\mathcal{I}_q(\mathfrak{M})$.

Before we prove that $\mathcal{I}_q(\overline{\mathfrak{M}})$ is a Gröbner basis of $\text{OS}_q(\mathfrak{M})$, we show that t_J^- reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$ for all $J \in \mathfrak{M}$.

Lemma 3.11. *Let $J, K \subseteq S$ be two dependent sets such that $J \cap K$ is independent. Then for all $l \in J \cup K$ the set $(J \cup K) \setminus \{l\}$ is dependent.*

Proof. Let $l \in J \cup K$. If there is a circuit contained in $J \setminus \{l\}$ or $K \setminus \{l\}$ then it is contained in $(J \cup K) \setminus \{l\}$. On the other hand, if $C \subseteq J$, $D \subseteq K$ are circuits containing l then $C \neq D$ since $J \cap K$ is independent. Hence by the circuit axiom there is a circuit contained in $(C \cup D) \setminus \{l\}$. $\square$

Proposition 3.12. *For any dependent set J , t_J^- reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$.*

Proof. We proceed by induction with respect to deglex order.

Let J be a dependent set. Recall that $\#J \geq 3$. If $J \in \overline{\mathfrak{M}}$ then t_J^- reduces to zero. In particular, this holds if J is the smallest dependent set with respect to deglex order.

Assume now that $J \notin \overline{\mathfrak{M}}$. Then there exists $K \in \overline{\mathfrak{M}}$ such that $K <_{\text{dlex}} J$ and $K \setminus \{k_1\} \subseteq J \setminus \{j_1\}$. In particular, K is dependent. We now distinguish several cases according to the relations between k_1 and the elements of J .

If $k_1 < j_1$ then let $L = \{k_1\} \cup (J \setminus \{j_1\})$. In this case $L <_{\text{dlex}} J$, $K \subset L$ and $K \setminus \{k_1\} \subseteq L \setminus \{l_1\}$. Hence t_L^- reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$ by induction hypothesis. If $L \cap J$ is dependent then $t_{L \cap J}^-$ reduces to zero by induction hypothesis and hence t_J^- reduces to zero by Lemma 3.9. Assume now that $L \cap J$ is independent. Then, by Lemma 3.11, $(J \cup L) \setminus \{j_i\}$ is dependent for all $i \in \{2, 3, \dots, \#J\}$ and is smaller than J with respect to $<_{\text{dlex}}$. We conclude from Lemma 3.2 that

$$0 = t_J^- - t_L^- + \sum_{i=2}^{\#J} (-1)^i t_{(J \cup L) \setminus \{j_i\}}^-$$

in A and hence t_J^- reduces to zero by induction hypothesis.

If $k_1 = j_1$ then t_J^- reduces to zero by using that t_K^- reduces to zero and by repeatedly applying Lemma 3.9.

If $j_1 < k_1 < j_2$ then $\#K < \#J$ since $K <_{\text{dlex}} J$. Since t_K^- reduces to zero, by repeatedly applying Lemma 3.9 we obtain a dependent set $L \subseteq S$ such that $\#L = \#J - 1$, $l_1 = k_1$, $L \setminus \{l_1\} \subseteq J \setminus \{j_1\}$, and t_L^- reduces to zero. There are two cases: $j_{\#J} \notin L$ or $j_2 \notin L$. If $J \cap L$ is dependent then t_J^- reduces to zero by induction hypothesis and by Lemma 3.9. If $J \cap L$ is independent then $(J \cup L) \setminus \{s\}$ is dependent for all $s \in J \cap L$ by Lemma 3.11. Now if $j_{\#J} \notin L$ then t_J^- reduces to zero by induction hypothesis and by Lemma 3.3. Observe that in Lemma 3.3 t_J^- and $t_L^- t_{j_{\#J}}$ are the summands with the largest leading term. Similarly, if $j_2 \notin L$ then t_J^- reduces to zero by induction hypothesis and by Lemma 3.4.

Finally, assume that $j_2 = k_1$ or $j_2 < k_1$. By repeatedly applying Lemma 3.9 we obtain that the set $L = \{s \in J \mid k_1 = s \text{ or } k_1 < s\}$ is dependent and $L <_{\text{dlex}} J$. If $l_1 \in J$ then t_J^- reduces to zero by Lemma 3.9. If $l_1 \notin J$ and $J \cap L$ is dependent then again J reduces to zero by induction hypothesis and Lemma 3.9. In the last case, if $J \cap L$ is independent then $(J \cup L) \setminus \{s\}$ is dependent for all $s \in L$, $l_1 < s$. In this case Lemma 3.2 applied to $J \setminus L$ and

L implies that

$$t_{J \setminus L}^- t_L^- = t_J^- + \sum_{i=2}^{\#L} (-1)^{i-1} t_{(J \cup L) \setminus \{l_i\}}^-$$

and the leading term on both sides of the equation is the leading term of t_J^- . Thus t_J^- reduces to zero by induction hypothesis. $\square$

4. INDEPENDENCE STATEMENT AND PROOFS

We first show that $I_q(\mathfrak{M})$ is independent of the chosen total order of S .

Proposition 4.1. *For any set system $\mathfrak{M}$ over S , the ideal $I_q(\mathfrak{M})$ and hence the algebra $\text{OS}_q(\mathfrak{M})$ are independent of the total order on S .*

Proof. We have to show that for any two total orders on S the defining ideals of $\text{OS}_q(\mathfrak{M})$ coincide. Relations (1.1), (1.2) are obviously independent of the chosen total order. Relations (1.2) can be used to reformulate a defining relation in (1.3) in terms of another order. We may simplify the problem by looking at orders $<$, $\ll$ which differ by exchanging two neighboring elements $a, b \in S$ with $a < b$, that is, $r \ll s$ for $r, s \in S$ if and only if either $(r, s) = (b, a)$ or $r < s$, $(r, s) \neq (a, b)$. We write $\ell_J^<(I)$ and $t_{J \setminus I}^<$ and similarly for $\ll$ to indicate the dependency on the order.

Let $J \in \mathfrak{M}$, $I \subseteq J$ with $2 \nmid \#I$. If $a \notin J$ or $b \notin J$ then

$$(-1)^{\ell_J^{\ll}(I)} = (-1)^{\ell_J^<(I)}, \quad (-q)^{(\#I-1)/2} t_{J \setminus I}^{\ll} = (-q)^{(\#I-1)/2} t_{J \setminus I}^<$$

and hence (1.3) takes the same form with respect to $<$ and $\ll$. It remains to consider the case when $a, b \in J$. We prove that in this case the defining relations differ by a sign. Then the proof of the proposition is completed.

Assume that $a \in I$, $b \in J \setminus I$ or $a \in J \setminus I$, $b \in I$. Then

$$(-1)^{\ell_J^{\ll}(I)} = -(-1)^{\ell_J^<(I)}, \quad t_{J \setminus I}^{\ll} = t_{J \setminus I}^<$$

and therefore $(-1)^{\ell_J^{\ll}(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}^{\ll} = -(-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}^<$. Assume now that $a, b \in J \setminus I$ and let $I' = I \cup \{a, b\}$. Then

$$\begin{aligned} \ell_J^{\ll}(I) &= \ell_J^<(I), \quad \ell_J^{\ll}(I') = \ell_J^<(I'), \quad t_{J \setminus I'}^{\ll} = t_{J \setminus I'}^<, \\ t_{J \setminus I}^{\ll} &= t_{j_1} \cdots (t_b t_a) \cdots t_{j_{\#J}} = t_{j_1} \cdots (-t_a t_b + 2q) \cdots t_{j_{\#J}} = -t_{J \setminus I}^< + 2q t_{J \setminus I'}^<, \end{aligned}$$

and $(-1)^{\ell_J^<(I)} = (-1)^{\ell_J^<(I')}$. Hence

$$\begin{aligned} &(-1)^{\ell_J^{\ll}(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}^{\ll} + (-1)^{\ell_J^{\ll}(I')} (-q)^{(\#I'-1)/2} t_{J \setminus I'}^{\ll} \\ &= (-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} (-t_{J \setminus I}^< + 2q t_{J \setminus I'}^<) + (-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} (-q) t_{J \setminus I'}^< \\ &= -(-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}^< + q (-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} t_{J \setminus I'}^< \\ &= -(-1)^{\ell_J^<(I)} (-q)^{(\#I-1)/2} t_{J \setminus I}^< - (-1)^{\ell_J^<(I')} (-q)^{(\#I'-1)/2} t_{J \setminus I'}^<. \end{aligned}$$

This is what we wanted to show. $\square$

Next we provide the proof of the main result.

Proof of Theorem 1.1. **Claim:** If $\mathfrak{M}$ is a loopless matroid without parallel elements then $I_q(\overline{\mathfrak{M}}) = I_q(\mathfrak{M})$.

◁ From Proposition 3.12 we deduce that $t_J^- \in I_q(\overline{\mathfrak{M}})$ for all circuits J . Hence $I_q(\mathfrak{M}) \subseteq I_q(\overline{\mathfrak{M}})$. To show equality it suffices to show that $t_J^- \in I_q(\mathfrak{M})$ for all dependent $J \subseteq S$. We prove the assertion by induction on the cardinality of the difference set of J and the circuit of largest cardinality contained in it. If the cardinality is 0 then J itself is a circuit and hence $t_J^- \in I_q(\mathfrak{M})$ by definition. If the cardinality is positive then there is an $s \in J$ such that $J \setminus \{s\}$ is dependent and by induction $t_{J \setminus \{s\}}^- \in I_q(\mathfrak{M})$. By Lemma 3.5(2) we can write t_J^- as an A linear combination of $t_{J \setminus \{s\}}^+$, $t_s t_{J \setminus \{s\}}^-$ and elements of $I_q(\emptyset) \subseteq I_q(\mathfrak{M})$. By Lemma 3.1 and since $t_{J \setminus \{s\}}^- \in I_q(\mathfrak{M})$ it follows that $t_{J \setminus \{s\}}^+ \in I_q(\mathfrak{M})$ and the assertion follows. ▷

We complete the proof by showing that $\mathcal{I}_q(\mathfrak{M})$ is a Gröbner basis of $I_q(\overline{\mathfrak{M}}) = I_q(\mathfrak{M})$. For this we verify that the conditions of Theorem 2.1 under the simplification provided by Lemma 2.2 are fulfilled.

First we have to find minimal generators of the modules $J_{f,g}$, where f, g are polynomials (1.1), (1.2) or t_J^- with $J \in \overline{\mathfrak{M}}$. According to Lemma 2.2 we can ignore generators $(\text{lm}(g) m, 1) \otimes (1, m \text{lm}(f))$ and $(1, \text{lm}(f) m) \otimes (m \text{lm}(g), 1)$ of $J_{f,g}$, where m is an arbitrary monomial. Further, since we do not fix an order on the Gröbner basis, we may restrict ourselves to generators of $J_{f,g}$ of the form $(1, m) \otimes (n, 1)$ and $(1, m) \otimes (1, n)$. Therefore the following cases have to be considered.

Case 1. $f = t_s^2 - q$, $g = t_s^2 - q$, $s \in S$.

The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_s, 1)$. Then $ft_s - t_sg$ is obviously zero.

Case 2. $f = t_s^2 - q$, $g = t_s t_r + t_r t_s - 2q$, $r, s \in S$, $r < s$.

The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_r, 1)$. The corresponding S -polynomial is

$$\begin{aligned} ft_r - t_sg &= t_s^2 t_r - q t_r - (t_s^2 t_r + t_s t_r t_s - 2q t_s) \\ &\searrow_{\mathcal{I}_q(\emptyset)} - q t_r - (-t_r t_s + 2q) t_s + 2q t_s \\ &\searrow_{\mathcal{I}_q(\emptyset)} - q t_r + t_r q = 0. \end{aligned}$$

Case 3. $f = t_s t_r + t_r t_s - 2q$, $g = t_r^2 - q$, $r, s \in S$, $r < s$.

The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_r, 1)$. The corresponding S -polynomial is

$$\begin{aligned} ft_r - t_sg &= (t_s t_r + t_r t_s - 2q) t_r - t_s (t_r^2 - q) \\ &\searrow_{\mathcal{I}_q(\emptyset)} t_r (-t_r t_s + 2q) - 2q t_r + q t_s \\ &\searrow_{\mathcal{I}_q(\emptyset)} - q t_s + q t_s = 0. \end{aligned}$$

Case 4. $f = t_s^2 - q$, $g = t_J^-$, $J \in \overline{\mathfrak{M}}$, $s = j_2$.

Let $K = J \setminus \{j_1, j_2\}$. The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_K, 1)$. The corresponding S -polynomial is

$$ft_K - t_sg = (t_s^2 - q) t_K - t_s t_J^-.$$

By Lemma 3.5 the expression $t_s t_J^-$ reduces to t_J^+ modulo $\mathcal{I}_q(\emptyset)$. In this reduction the leading term $t_s^2 t_K$ of $t_s t_J^-$ has to be reduced at one moment to $q t_K$. Therefore $t_s t_J^- - (t_s^2 - q) t_K$ also reduces to t_J^+ modulo $\mathcal{I}_q(\emptyset)$. Thus $t_s t_J^- - (t_s^2 - q) t_K$ reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$ by Lemma 3.8.

Case 5. $f = t_J^-$, $g = t_s^2 - q$, $J \in \overline{\mathfrak{M}}$, $s = j_{\#J}$.

Let $K = J \setminus \{j_1, j_{\#J}\}$. The remaining generator of $J_{f,g}$ is $(1, t_K) \otimes (t_s, 1)$. The corresponding S -polynomial is

$$f t_s - t_K g = t_J^- t_s - t_K (t_s^2 - q).$$

By Lemma 3.7 the expression $t_J^- t_s$ reduces to $(-1)^{\#J+1} t_J^+$ modulo $\mathcal{I}_q(\emptyset)$. In this reduction the leading term $t_K t_s^2$ of $t_J^- t_s$ has to be reduced at one moment to $q t_K$. Therefore $t_J^- t_s - t_K (t_s^2 - q)$ also reduces to $(-1)^{\#J+1} t_J^+$ modulo $\mathcal{I}_q(\emptyset)$. Thus $t_J^- t_s - t_K (t_s^2 - q)$ reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$ by Lemma 3.8.

Case 6. $f = t_s t_r + t_r t_s - 2q$, $g = t_r t_p + t_p t_r - 2q$, $p, r, s \in S$, $p < r < s$.

The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_p, 1)$. The corresponding S -polynomial is

$$\begin{aligned} f t_p - t_s g &= (t_s t_r + t_r t_s - 2q) t_p - t_s (t_r t_p + t_p t_r - 2q) \\ &= t_r t_s t_p - 2q t_p - t_s t_p t_r + 2q t_s \\ &\searrow^{\mathcal{I}_q(\emptyset)} t_r (-t_p t_s + 2q) - (-t_p t_s + 2q) t_r - 2q t_p + 2q t_s \\ &\searrow^{\mathcal{I}_q(\emptyset)} - (-t_p t_r + 2q) t_s + t_p (-t_r t_s + 2q) - 2q t_p + 2q t_s = 0. \end{aligned}$$

Case 7. $f = t_s t_r + t_r t_s - 2q$, $g = t_J^-$, $r, s \in S$, $r < s$, $J \in \overline{\mathfrak{M}}$, $r = j_2$.

Let $K = J \setminus \{j_1, j_2\}$. The remaining generator of $J_{f,g}$ is $(1, t_s) \otimes (t_K, 1)$. The corresponding S -polynomial is

$$f t_K - t_s g = (t_s t_r + t_r t_s - 2q) t_K - t_s t_J^-.$$

By Lemma 3.5 the expression $t_s t_J^-$ reduces to $\pm t_{J \cup \{s\}}^- + t_J^+$ modulo $\mathcal{I}_q(\emptyset)$. In this reduction the leading term $t_s t_r t_K$ of $t_s t_J^-$ has to be reduced at one moment to $(-t_r t_s + 2q) t_K$. Therefore $t_s t_J^- - (t_s t_r + t_r t_s - 2q) t_K$ also reduces to $\pm t_{J \cup \{s\}}^- + t_J^+$ modulo $\mathcal{I}_q(\emptyset)$. Since t_J^+ reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$ by Lemma 3.8, it suffices to prove that $t_{J \cup \{s\}}^-$ reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$. The latter holds for $s \leq j_2$ and for $j_{\#J} \leq s$ by Lemma 3.9 and for $j_2 < s < j_{\#J}$ by (GC).

Case 8. $f = t_J^-$, $g = t_s t_r + t_r t_s - 2q$, $r, s \in S$, $r < s$, $J \in \overline{\mathfrak{M}}$, $s = j_{\#J}$.

Let $K = J \setminus \{j_1, s\}$. The remaining generator of $J_{f,g}$ is $(1, t_K) \otimes (t_r, 1)$. The corresponding S -polynomial is

$$f t_r - t_K g = t_J^- t_r - t_K (t_s t_r + t_r t_s - 2q).$$

The proof is similar to the one in Case 7 and uses Lemma 3.7.

Case 9. $f = t_J^-$, $g = t_K^-$, $J \setminus \{j_1\} \subseteq K \setminus \{k_1\}$, $J, K \in \overline{\mathfrak{M}}$, $J \neq K$. This case does not appear by Condition (GC) on the elements of $\overline{\mathfrak{M}}$.

Case 10. $f = t_J^-$, $g = t_K^-$, $J, K \in \overline{\mathfrak{M}}$, there exists $i \in \{3, \dots, \#J\}$ such that $j_i = k_2$, $j_{i+1} = k_3, \dots, j_{\#J} = k_{\#J-i+2}$, $\#J - i + 2 < \#K$.

Let $n = \#J - i + 2$,

$$J' = J \setminus \{j_1\}, \quad K' = K \setminus \{k_1\}, \quad L = \{j \in J' \mid j < k_1\}, \quad M = \{j \in J' \setminus K' \mid k_1 < j\}.$$

Thus the sets L , M and $J' \cap K'$ are pairwise disjoint and their union is J' (if $k_1 \notin J'$) or $J' \setminus \{k_1\}$ (if $k_1 \in J'$). We have to show that

$$(4.1) \quad t_J^- t_{K' \setminus J'} - t_{J' \setminus K'} t_K^- \searrow^{\mathcal{I}_q(\overline{\mathfrak{M}})} 0.$$

We will proceed in several steps, and at some moment we will have to distinguish the cases $k_1 \in J'$ and $k_1 \notin J'$.

Using Lemma 3.1 and Lemma 3.8 we observe first that

$$(4.2) \quad t_J^- t_{K' \setminus J'} = t_{J \cup K'}^- + \text{lower terms which reduce to zero modulo } \mathcal{I}_q(\overline{\mathfrak{M}}).$$

Further, by applying Lemma 3.5(2) and Lemma 3.8 we obtain that

$$t_M t_K^- = t_{M \cup K}^- + \text{lower terms which reduce to zero modulo } \mathcal{I}_q(\overline{\mathfrak{M}}).$$

In particular, if $j_1 = k_1$ then $M \cup K = J' \cup K = J \cup K'$ and hence (4.1) holds.

Assume now that $k_1 \in J'$. Then $t_{J' \setminus K'} = t_L t_{k_1} t_M$. Lemma 3.5(1) gives that

$$t_{k_1} t_{M \cup K}^- = t_{M \cup K}^+ + \text{lower terms which reduce to zero modulo } \mathcal{I}_q(\overline{\mathfrak{M}})$$

and Equation (3.1) implies that

$$t_L t_{M \cup K}^+ = t_{L \cup M \cup K}^+ + \text{lower terms which reduce to zero modulo } \mathcal{I}_q(\overline{\mathfrak{M}}).$$

Now, $L \cup M \cup K = (J \cup K) \setminus \{j_1\}$ and

$$(4.3) \quad t_{J \cup K'}^- = t_{J \cup K}^- = t_{(J \cup K) \setminus \{j_1\}}^+ - t_{j_1} t_{(J \cup K) \setminus \{j_1\}}^-$$

by Equation (3.2). Since $K \subseteq (J \cup K) \setminus \{j_1\}$, Proposition 3.12 yields that $t_{(J \cup K) \setminus \{j_1\}}^-$ reduces to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$. Thus we conclude from (4.2) and (4.3) that (4.1) holds.

Finally, assume that $k_1 \notin J$. Then $t_{J' \setminus K'} = t_L t_M$. Further,

$$t_L t_{M \cup K}^- = t_{\{j_1\} \cup L}^- t_{M \cup K}^- + \text{lower terms which reduce to zero modulo } \mathcal{I}_q(\overline{\mathfrak{M}})$$

by definition of $t_{\{j_1\} \cup L}^-$, and hence

$$(4.4) \quad t_L t_{M \cup K}^- = \sum_{m=1}^{\#(M \cup K)} (-1)^{m-1} t_{(J \cup K) \setminus \{(M \cup K)_m\}}^- + \text{lower terms which reduce to zero}$$

by Lemma 3.2, where $(M \cup K)_m$ is the m th element of $M \cup K$. The summand containing the leading term of the last expression is $t_{(J \cup K) \setminus \{k_1\}}^-$ since $(M \cup K)_1 = k_1$. Since $J \cap K$ is independent by the assumptions $J \cap K = J \cap K' \subseteq K'$ and $K \in \overline{\mathfrak{M}}$, Lemma 3.11 and Proposition 3.12 imply that all other summands in (4.4) reduce to zero modulo $\mathcal{I}_q(\overline{\mathfrak{M}})$. Thus (4.1) holds in this case. $\square$

It remains to provide the proof of Corollary 1.4.

Proof of Corollary 1.4. **Claim:** There is a total order on S for which all Gröbner circuits are of size 3.

◁ We recall the characterization of supersolvable matroids given in [BZ91, Thm. 2.8 (5)]. There it is shown that for a supersolvable matroid $\mathfrak{M}$ on ground set S the set S can be partitioned into subsets $S = S_1 \cup \cdots \cup S_f$ such that for any $1 \leq h \leq f$ and two elements $x, y \in S_h$ there is an $1 \leq g < h$ and $z \in S_g$ such that $\{x, y, z\} \in \mathfrak{M}$ is a circuit. Now we choose a total order on S such that for $1 \leq g < h \leq f$ all elements from S_g come before S_h . Assume that J is a Gröbner circuit in this order. If $J \cap S_h \geq 2$ for some $1 \leq h \leq f$ then by [BZ91, Thm. 2.8 (5)] for any two elements $x, y \in J \cap S_h$ there is a circuit K of size 3 such that $K \setminus \{k_1\} = \{x, y\}$. By choosing two elements $\{x, y\}$ from $J \cap S_h$ for which $\{x, y\} \sqsubseteq J$ we can choose K such that $K \leq_{\text{dlex}} J$ and $K \setminus \{k_1\} \sqsubseteq J \setminus \{j_1\}$. From this it follows by (GC) that $J = K$. ▷

Now if all Gröbner circuits are of size 3 then the Gröbner basis from Corollary 1.3 is quadratic. Hence by well known facts (see [Frö99, Sec. 4]) it follows that $\text{OS}_q(\mathfrak{M})$ is Koszul. $\square$

ACKNOWLEDGEMENTS

We thank Graham Denham, Bernd Sturmfels and Alexander Suciu for useful comments on an earlier version of this paper.

REFERENCES

- [BZ91] A. Björner and G.M. Ziegler, *Broken circuit complexes: Factorizations and generalizations*, J. Comb. Theory, Ser. B **51** (1991), 96–126.
- [CF05] R. Cordovil and D. Forge, *Gröbner and diagonal bases in Orlik-Solomon type algebras*, Cubo **7** (2005), 1–20.
- [Frö99] R. Fröberg, *Koszul algebras*, Advances in commutative ring theory. Proceedings of the 3rd international conference, Fez, Morocco (New York), Lect. Notes Pure Appl. Math., vol. 205, Marcel Dekker, 1999, pp. 337–350.
- [Li02] H. Li, *Noncommutative Gröbner bases and filtered-graded transfer*, Lect. Notes in Math., vol. 1795, Springer-Verlag, Berlin, Heidelberg, 2002.
- [Mor94] T. Mora, *An introduction to commutative and noncommutative Gröbner bases*, Theor. Comput. Sci. **134** (1994), 131–173.
- [Orl07] P. Orlik, *Lectures on arrangements: combinatorics and topology*, Algebraic combinatorics (Heidelberg) (G. Fløystad, ed.), Springer, 2007, pp. 1–79.
- [OS80] P. Orlik and L. Solomon, *Combinatorics and topology of complements of hyperplanes*, Invent. Math. **56** (1980), 167–189.
- [Oxl06] J. G. Oxley, *Matroid theory*, Oxford Graduate Texts in Mathematics, vol. 3, Oxford University Press, Oxford, 2006.
- [Pee03] I. Peeva, *Hyperplane arrangements and linear strands in resolutions*, Trans. Amer. Math. Soc. **355** (2003), 609–618.
- [SY97] B. Shelton and S. Yuzvinsky, *Koszul algebras from graphs and hyperplane arrangements*, J. Lond. Math. Soc., II. Ser. **56** (1997), 477–490.
- [Wel76] D.J.A. Welsh, *Matroid theory*, L.M.S. Monographs, vol. 8, Academic Press, London, 1976.
- [Yuz01] S. Yuzvinsky, *Orlik-Solomon algebras in algebra and topology*, Russ. Math. Surv. **56** (2001), 293–364.

FACHBEREICH MATHEMATIK UND INFORMATIK, PHILIPPS-UNIVERSITÄT MARBURG, 35032 MARBURG, GERMANY

E-mail address: `heckenberger@mathematik.uni-marburg.de`

E-mail address: `welker@mathematik.uni-marburg.de`